

**<20% TBSA BURN CHECKLIST**

- ☐ Burn Consult order placed
- ☐ Pictures of burns in chart (can be done by burn team during daytime)
- ☐ **Perform mapping on Lund-Browder Chart**  
(see reverse, perform after debridement, this will serve as initial and final mapping if discharged)
- ☐ If appropriate for discharge, burn-specific discharge instructions and clinic message sent per Burn ED Discharge Guideline
- ☐ If requires admission, admit to Burn service as primary team with **Burn A attending** as admitting attending
- ☐ If requires admission, place PT/OT consult

**>20% TBSA → transport to TBICU before debridement**

**<20% TBSA → Debride in trauma bay before final disposition**

**Indications for CT scan:**

- Fall >6 feet
- Explosion
- Major traumatic mechanism

- If 7a-3p, coordinate debridement with Burn Dressing Team
- If 3p-12a, trauma chief and TBICU resident will be responsible for debridement
- If 12a-7a, patient will be debrided at ~9AM by Burn Dressing Team

**>20% TBSA BURN CHECKLIST****IMMEDIATE TRAUMA BAY INTERVENTIONS**

- ☐ Start resuscitation sheet
- ☐ Record ED arrival time on resuscitation sheet
- ☐ Start crystalloid **immediately at minimum rate 300 mL/hr**
- ☐ Insert temperature-sensing foley
- ☐ If temperature <37°C, place bair hugger on patient

**INITIAL MAPPING AND FLUID CALCULATION (IN TRAUMA BAY, BEFORE DEBRIDEMENT)**

- ☐ Perform initial mapping using Lund-Browder chart on reverse side
- ☐ Calculate initial fluid resuscitation rate **(2 mL/kg/%TBSA ÷ 16)**
- ☐ Record time of initial mapping and calculated fluid resuscitation on resuscitation sheet (Goal = 30 minute from arrival)
- ☐ Cover patient with ¾ impervious drape and blankets until initial debridement occurs

**TBICU ADMISSION**

- ☐ Expedite TBICU admission (Goal = 1 hour from arrival)
- ☐ Burn service as primary team with Burn A attending as admitting attending
- ☐ Record TBICU arrival time on resuscitation sheet
- ☐ If >30% TBSA, notify burn attending
- ☐ If >30% TBSA, insert Dobhoff tube and initiate tube feeds at 30 kcal/kg/24hr
- ☐ **Coordinate initial debridement**  
(can be delayed up to 8hrs, ensure order for polysporin or bacitracin is placed)
- ☐ Give TBICU nurses 2 hours to admit patient before starting debridement. Debridement should not occur within 2 hours of shift change.
- ☐ Touch base with ANM prior to beginning debridement to ensure adequate resources
- ☐ Place PT/OT Consult

Signature: \_\_\_\_\_ Date: \_\_\_\_\_

